

Important Safety Information on NARDIL (phenelzine sulfate) Shortage

Updated on 2021/02/17

2020/11/23

Audience

Healthcare professionals including psychiatrists, general practitioners, pharmacists, and nurses.

Key messages

- **An ongoing global shortage of NARDIL (phenelzine sulfate), a monoamine oxidase inhibitor (MAOI) indicated for the treatment of certain types of depression, is expected to continue due to a lack of the active ingredient.**
- **Healthcare professionals are advised to:**
 - **Ensure no new patients are started on NARDIL.**
 - **Initiate switching patients taking NARDIL to an alternative treatment.**
 - **Avoid abrupt discontinuation of NARDIL treatment to prevent withdrawal syndrome, which can be serious. Treatment discontinuation should be gradual, and a `wash out` period of at least 10 days is necessary after discontinuation of NARDIL and before the start of the new treatment.**
 - **Monitor patients closely when taking patients off of NARDIL.**
- **Erfa Canada 2012 Inc., in collaboration with Health Canada, is working to help minimize the impact of this shortage on Canadians.**

What is the issue?

Due to an ongoing global shortage of the active ingredient, the availability of NARDIL (phenelzine sulfate) in Canada, manufactured by ERFA Canada 2012 Inc., has decreased and the

ERFA Canada 2012 Inc.

110-8250 boul. Décarie Montréal, QC, Canada, H4P 2P5
Tel: (514) 931-3133 Toll free: 1-888-922-3133 Fax: (514) 931-7330
www.eci2012.net

resupply date is unknown at this point.

Phenelzine is currently listed on the [Tier 3 list](#) of shortages. Tier 3 shortages are those with the greatest impact on supply and drugs on this list are in high demand or currently in shortage. Health Canada works in collaboration with the provinces and territories, industry and stakeholders to mitigate the impact of Tier 3 shortages on patients and the health care system. Additional information and the latest updates regarding the shortage of NARDIL are available at: drugshortagescanada.ca.

Products affected

NARDIL, Phenelzine Sulfate Tablets USP, 15 mg DIN 00476552.

Background information

NARDIL is a potent monoamine oxidase inhibitor (MAOI) indicated for the treatment of "atypical", "nonendogenous" or "neurotic" depression. NARDIL is indicated for patients who have failed to respond to other drugs more commonly used for these conditions.

Due to a global shortage, NARDIL has been on backorder since March 2020. In March 2020, ERFA Canada 2012 Inc. sent a backorder notification to Health Canada, and a letter to all Canadian pharmacies outlining the importance of informing NARDIL patients and their physicians about the actual shortage and advising them to consider switching to alternative treatments. On the 24th of June 2020, ERFA Canada 2012 Inc. in coordination with Health Canada and other Canadian stakeholders, issued a Risk Communication outlining the recommendations for healthcare professionals to not start new patients on Nardil and to switch the existing patients to other alternative treatments. Current shortage information is available at www.drugshortagescanada.ca.

Information for consumers

NARDIL (phenelzine sulfate) is used to treat depression where anxiety or fear is the main symptom and treatment with other drugs has failed.

There is a global shortage of NARDIL and based on current available information, the Canadian product manufacturer cannot guarantee a steady supply in the future. All patients who are treated with NARDIL should consult their healthcare professional as soon as possible in order to discuss alternative treatment options.

Before patients stop taking NARDIL, they need to talk to their healthcare professional. Patients should discuss with their healthcare professional how to safely discontinue NARDIL to minimize discontinuation symptoms and understand what symptoms require medical attention. Patients who suddenly stop taking NARDIL can experience serious side effects within one to three days including nightmares, agitation, psychosis (seeing or hearing things that are not there, or

ERFA Canada 2012 Inc.

110-8250 boul. Décarie Montréal, QC, Canada, H4P 2P5
Tel: (514) 931-3133 Toll free: 1-888-922-3133 Fax: (514) 931-7330
www.eci2012.net

believing things which are not true) and convulsions.

Patients are advised to contact their health care professional and seek medical advice as soon as possible should they experience these side effects or an exacerbation of their underlying disease. Patients should also discuss any questions or concerns with their healthcare professional.

Information for healthcare professionals

Abrupt withdrawal of NARDIL can be associated with an uncommon withdrawal syndrome. Signs and symptoms of this syndrome generally commence 24 to 72 hours after drug discontinuation and may range from vivid nightmares with agitation to frank psychosis and convulsions. This syndrome generally responds to reinstatement of low-dose NARDIL therapy followed by cautious downward titration and discontinuation.

Healthcare professionals are advised to:

- Ensure no new patients are started on NARDIL.
- Initiate switching patients taking NARDIL to an alternative treatment.
- Avoid abrupt discontinuation of NARDIL treatment to prevent withdrawal syndrome, which can be serious. NARDIL dose should be tapered off gradually for 3 to 5 weeks, and a 'wash out' period of at least 10 days is necessary after discontinuation of NARDIL and before the start of the new treatment.
- Monitor patients closely when taking patients off NARDIL.

Action taken by Health Canada

Health Canada and health care system stakeholders are working with ERFA Canada 2012 Inc. to manage the supply disruption of NARDIL. Health Canada has communicated important safety information to healthcare professionals and Canadians via the [Recalls and Safety Alerts Database](#) on the Healthy Canadians Web Site.

Report health or safety concerns

Managing marketed health product-related side effects depends on healthcare professionals and consumers reporting them. Any serious or unexpected side effects in patients receiving NARDIL should be reported to Health Canada.

ERFA Canada 2012 Inc.

110-8250 boul. Décarie Montréal, QC, Canada, H4P 2P5
Tel: (514) 931-3133 Toll free: 1-888-922-3133 Fax: (514) 931-7330
www.eci2012.net

Erfa Canada 2012 Inc.

8250 BLVD DECARIE, SUITE 110

Montreal, Quebec

H4P 2P5

Tel: 1-514-931-3133

Fax: 1-514-931-7330

Email: psq@eci2012.net

You can report any suspected adverse reactions associated with the use of health products to Health Canada by:

- Calling toll-free at 1-866-234-2345; or
- Visiting MedEffect Canada's Web page on [Adverse Reaction Reporting](https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html) (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax.

For other health product inquiries related to this communication, contact Health Canada at:

Marketed Health Products Directorate

E-mail: mailto:mhpd_dpdc.public@hc-sc.gc.ca

Telephone: 613-954-6522

Fax: 613-952-7738

Original signed by

Mjellma Hatipi

Medical Director

Erfa Canada 2012 Inc.

ERFA Canada 2012 Inc.

110-8250 boul. Décarie Montréal, QC, Canada, H4P 2P5

Tel: (514) 931-3133 Toll free: 1-888-922-3133 Fax: (514) 931-7330

www.eci2012.net